

Medical Science

To Cite:

Rogowska Z, Omid S, Piątkiewicz M, Mazepa W, Olędzka M, Pruszkowska K, Olędzka B, Rzepko A. Drug reaction with eosinophilia and systemic symptoms (DRESS Syndrome) - A comprehensive review of clinical features, optimized diagnosis and management strategies. *Medical Science* 2026; 30: e174ms3912 doi: <https://doi.org/10.54905/disssi.v30i174.e174ms3912>

Authors' Affiliation:

¹Faculty of Medicine, Medical University of Warsaw, Żwirki I Wigury 61, 02-091 Warsaw, Poland

²Maria Skłodowska-Curie Medical University of Warsaw, Aleja Solidarności 12 Street, 03-411 Warsaw, Poland

ORCID:

Zofia Rogowska: 0009-0001-7605-2216
Sara Omid: 0009-0009-3462-9222
Michał Piątkiewicz: 0009-0005-5182-3407
Wiktoria Mazepa: 0009-0006-9719-6733
Małgorzata Olędzka: 0009-0002-3952-0380
Kaja Pruszkowska: 0009-0006-6541-2868
Barbara Olędzka: 0009-0005-7785-8659
Aleksandra Rzepko: 0009-0006-1459-5027

*Corresponding author:

Zofia Rogowska,
Faculty of Medicine, Medical University of Warsaw, Żwirki I Wigury 61, 02-091 Warsaw, Poland
e-mail: zrogowska@op.pl
ORCID: 0009-0001-7605-2216

Peer-Review History

Received: 21 March 2026
Reviewed & Revised: 03/April/2026 to 09/August/2026
Accepted: 18 August 2026
Published: 29 August 2026

Peer-review Method

External peer-review was done through double-blind method.

Medical Science
pISSN 2321-7359; eISSN 2321-7367



© The Author(s) 2026. Open Access. This article is licensed under a Creative Commons Attribution License 4.0 (CC BY 4.0), which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons license, and indicate if changes were made. To view a copy of this license, visit <http://creativecommons.org/licenses/by/4.0/>.

Drug reaction with eosinophilia and systemic symptoms (DRESS Syndrome) - A comprehensive review of clinical features, optimized diagnosis and management strategies

Zofia Rogowska^{1*}, Sara Omid¹, Michał Piątkiewicz¹, Wiktoria Mazepa¹, Małgorzata Olędzka², Kaja Pruszkowska¹, Barbara Olędzka¹, Aleksandra Rzepko¹

ABSTRACT

Drug reaction with eosinophilia and systemic symptoms, also known as drug-induced hypersensitivity syndrome (DRESS/ DIHS), represents a rare, severe adverse drug reaction. The syndrome typically manifests with rash, fever, facial edema, and multiorgan involvement and requires a prompt withdrawal of the culprit drug. This review summarises existing knowledge on DRESS, including pathophysiology, clinical presentation, diagnostic approaches and treatment options.

Keywords: Drug hypersensitivity syndrome, Drug eruption, DRESS, Drug-related side effects and adverse reactions, Eosinophilia

1. INTRODUCTION


DRESS syndrome, first described in 1996 by Bocquet, is a rare, severe, and potentially life-threatening cutaneous adverse hypersensitivity reaction (Bocquet et al., 1996; Peter et al., 2017). The Japanese research group later proposed the term “Drug-induced hypersensitivity syndrome” (DiHS), which is often used interchangeably to describe the same syndrome. The disease usually begins 2-8 weeks after administration of the causative drug. It is most often associated with the use of anticonvulsants, antibiotics, allopurinol, and non-steroidal anti-inflammatory agents (NSAIDs), although many other drugs may also be involved in the pathogenesis.

The clinical presentation of the disease is progressive. DRESS is characterized by generalized skin rash with facial swelling, fever, lymphadenopathy, hematologic abnormalities, particularly eosinophilia and lymphocytosis, and more or less severe involvement of visceral organs, such as hepatitis, nephritis, pneumonia, and myocarditis. The pathophysiology of this heterogeneous syndrome is complex, multifactorial, and not fully understood (Kroshinsky et al., 2024). Viral reactivation of HHV-6/7, EBV, and CMV is a significant aspect of disease presentation. This viral

reactivation may cause a prolonged antiviral response that can lead to the development of systemic effects (Hama et al., 2022; Miyagawa et al., 2021). In the treatment of DRESS syndrome, it is important to discontinue the drug causing the hypersensitivity reaction. In addition to prompt withdrawal of all suspected causative drugs, treatment includes administration of glucocorticosteroids or other immunosuppressive agents, as indicated.

2. REVIEW METHODS

To write this review, we used the scientific database PubMed. We focused on recent, relevant publications written in English. The search focused on studies addressing the clinical features, pathogenesis, diagnosis, and management strategies of DRESS syndrome. We selected studies in an advanced search strategy using keywords related to the subject, such as „Drug hypersensitivity syndrome“, „DRESS“, „Eosinophilia“, „Drug eruption“, or „Drug-related side effects and adverse reactions“. To prepare articles for this study, we conducted research from December 2025 to February 2026. The selection procedures are shown in Figure 1.



*n- number of articles

Figure 1. Flow chart

3. RESULTS AND DISCUSSION

Epidemiology

The true incidence of DRESS varies depending on the drug involved and the patient's immune status and is likely underestimated due to underdiagnosis and underreporting. It's reported incidence lies between 2 and 5 cases per million per year with estimated prevalence being approximately 2 cases per 100,000 population (Kroshinsky et al., 2024).

DRESS syndrome can occur at any age but is seen more often in adults, with age distribution varying by population and causative drug (Del Pozzo-Magaña et al., 2022; Rahnama-Moghadam et al., 2025; Toniato et al., 2021). A higher incidence of DRESS has been observed among Black individuals and women (Hiransuthikul et al., 2016; Wang and Mei, 2017; Kardaun et al., 2013). Despite appropriate treatment, the mortality rate remains between 3.8% and 10%. (Hiransuthikul et al., 2016).

Etiology

At least 60 medications have been associated with DRESS. Those most frequently implicated are aromatic anticonvulsants (phenytoin, carbamazepine, and phenobarbital), sulfonamides, sulfones (dapsone), nonsteroidal anti-inflammatory drugs (piroxicam, ibuprofen, and diclofenac), beta-lactam antibiotics, vancomycin, allopurinol, minocycline, and antiretrovirals. Criado et al., (2025) and Liang et al., (2024) reported that the FAERS database for over 20 years up to 2023 recorded 15,751 cases of DRESS, of which 1,043 were fatal. Study showed that the most common drugs causing DRESS were two antibiotics, while allopurinol was the agent most responsible for death cases. Table 1 describes the main medications listed as potential triggers for DRESS (Stirton et al., 2022; Criado et al., 2025; Calle et al., 2023).

Table 1. Summary of the most commonly associated drugs.

Drug Category	Drug Name
Anticonvulsants - Aromatic	Carbamazepine, phenytoine, lamotrigine, phenobarbital, oxcarbazepine
Anticonvulsants - Non-aromatic	Ethosuximide, sodium valproate or valproic acid, zonisamide, gabapentin, levetiracetam
Antibiotics	Sulfonamides, sulfamethoxazole-trimethoprim, sulfasalazine, dapsone, vancomycin, minocycline, ampicillin, ampicillin/sulbactam, azithromycin, ceftriaxone, cefepime, clindamycin, spiramycine, metronidazole, amoxicillin, amoxicillin/clavulanate, linezolid, clindamycin, levofloxacin, hydroxychloroquine, teicoplanin, voriconazole
Antituberculosis agents	rifampin, ethambutol, pyrazinamide, isoniazid, streptomycin
Analgesics	Diclofenac, celecoxib, ibuprofen, acetylsalicylic acid, piroxicam, metamizole, dextketoprofen
Antidepressant and antipsychotic	Bupropion, fluoxetine, olanzapine, quetiapine, desipramine, amitriptyline, clomipramine,
Antihypertensive	Amlodipine, captopril, enalapril, diltiazem, spironolactone
Antivirals	Telaprevir, boceprevir, abacavir, raltegravir, nevirapine, zalcitabine
Others	Allopurinol, omeprazole, esomeprazole, propylthiouracil, nivolumab, ipilimumab, Traditional Chinese Medicine, quinine, mexiletine, hydroxychloroquine, ranitidine, thiamine, vitamin B12, rivaroxaban

Pathophysiology

Pathophysiology of DRESS primarily involves a complex interaction between immunological mechanisms and genetic predisposition. The disease is mediated by T-cell activation and cytokine release (Criado et al., 2025). Moreover, certain human leukocyte antigens have been linked to different drug hypersensitivity reactions, which also include DRESS syndrome. Also, the relation between reactivation of latent viruses, such as HHV-6, plays a vital role in the development of the syndrome (Stirton et al., 2022; Criado et al., 2025; Calle et al., 2023). Additional predisposing factors likely exist but remain still unknown. It highlights the need to change this model as our understanding of DRESS pathogenesis evolves (Hung et al., 2005; Calle et al.,2023).

Clinical manifestations

DRESS syndrome typically develops within 2 weeks to 2 months after the initiation of the offending drug and is characterized by progressive multi-organ involvement, affecting the skin, hematologic system, and various internal organs (Stirton et al., 2022). DRESS syndrome typically begins with prodromal flu-like symptoms, including malaise, pruritus, pharyngitis, lymphadenopathy, and high fever (38–40 °C), which occurs in 75–100% of patients and often precedes the rash by several days. The clinical course of DRESS syndrome is usually gradual, with progressive involvement of multiple symptoms (Cacoub et al., 2011; Criado et al., 2025; Kardaun et al., 2013).

Cutaneous manifestations of DRESS are diverse and typically involve over 50% of the body surface area. The most characteristic early signs, present in up to 75% of patients, are periorbital and facial edema, often accompanied by micropustules. Facial edema may be a distinguishing feature from more mild forms of DRESS or maculopapular eruption (MPE). (Shiohara and Mizukawa, 2019) The rash in DRESS is often polymorphic (in around 85% of cases), including maculopapular, lichenoid, exfoliative, urticarial, purpuric, eczematous, and pustular lesions, with widespread erythema (Lehloenya et al., 2020; Kardaun et al., 2013)

Most patients develop a maculopapular exanthema that progresses in a craniocaudal, symmetrical manner, sparing the palms and soles (Shiohara and Mizukawa, 2019).

Mucosal involvement occurs in up to 56% of DRESS patients, usually affecting a single site such as the lips, pharynx, or tonsils, and may occasionally progress to erosions. It is typically mild and non-hemorrhagic, which helps to distinguish it from SJS/TEN (Eshki et al., 2009). In 20–30% of cases, erythema progresses to erythroderma. (Criado et al., 2025) Cutaneous manifestations may continue for several weeks to months even after the causative drug has been withdrawn (Cacoub et al., 2011).

Hematological abnormalities are common in DRESS, with atypical lymphocytosis often being one of the most frequent findings, though it can be easily overlooked. Eosinophilia, appearing later in 60–90% of patients across multiple studies (although the majority show greater than 80%), may take 1–2 weeks to develop and can persist even after liver enzymes have normalized (Stirton et al., 2022; Calle et al., 2023; Kardaun et al., 2013). Less commonly, DRESS may present with lymphopenia, leukopenia, thrombocytopenia, thrombocytosis, or pancytopenia, which are generally linked to a more severe prognosis (Stirton et al., 2022).

Systemic involvement of internal organs in DRESS is observed in up to 91% of patients. The most common manifestations of visceral organ involvement are hepatitis, nephritis, and pneumonia. Lymphadenopathy is seen in 50–75% of cases. Hepatic injury is common in patients with DRESS syndrome, occurring in up to 97% of cases. Typically it is characterised by an elevation of liver enzymes, which is usually mild, anicteric and may take months to normalize. The primary cause of mortality in DRESS is a rarely seen severe liver failure, occurring with extensive necrosis, coagulopathy and encephalopathy (Criado et al., 2025).

The second most commonly affected organ in DRESS is the kidney. Renal manifestations occur in up to 30% of DRESS cases. Most of the time, the injury is mild and resolves after discontinuation of the causative medication. It may present as moderate increases in creatinine, proteinuria, and eosinophils in the urinary sediment. However, sometimes renal impairment may be severe and manifest as interstitial nephritis, potentially progressing to end-stage renal disease. Patients at highest risk of kidney failure include the elderly, those with pre-existing renal disease, and cases of DRESS associated with allopurinol (Taweeseet et al., 2019).

The third most commonly affected organ in DRESS is the lung. Studies have shown that interstitial pneumonitis is the main pulmonary manifestation. Minocycline is the drug most frequently linked to pulmonary damage (Criado et al., 2025). Cardiac involvement in DRESS, typically manifests as eosinophilic myocarditis or pericarditis, which can have potentially fatal outcomes. Most commonly, it appears around 70 days after symptom onset and can occur even months after drug discontinuation. Heart failure in DRESS is characterized by chest pain, tachycardia, hypotension and dyspnea. It has typically been reported after administration of minocycline (Calle et al., 2023).

In addition to the more common organ involvement, patients with DRESS may occasionally develop pancreatitis, colitis, cholangitis, encephalitis or meningoencephalitis, hemophagocytic syndrome, and thyroiditis (Criado et al., 2025; Cacoub et al., 2011; Kardaun et al., 2013).

How to identify the causative DRESS drug?

Knowledge about the drug responsible for causing DRESS syndrome is crucial to stop disease progression and prevent re-exposure. It has been reported that intradermal testing can detect both immediate and delayed drug adverse reactions. Previously, specialists were divided on the use of those tests in severe reactions; however, with certain precautions, they are now widely considered a safe and useful tool for identifying the causative DRESS drug. The lymphocyte transformation test is another valuable, widely used diagnostic

tool for identifying delayed-type allergic reactions. It may help detect the causative drug by measuring drug-specific lymphocyte proliferation. When performed 4–8 weeks after the reaction, ideally within 6 months, it has a sensitivity of up to 73% and specificity of 85%. Moreover, it can be enhanced when combined with patch testing (Cabañas et al., 2018). It is important to note that the lymphocyte transformation test does not carry the risk of provoking a reaction through drug re-exposure. However, there is still a lack of complete standardization of this method (Kardaun et al., 2013).

When the lymphocyte transformation test is unavailable, patch testing is performed. Studies have shown its safety in immunocompetent patients (Cabañas et al., 2018). Patch testing should be used 4–6 weeks after the reaction. The positive predictive value of this method reaches up to 80% for carbamazepine but only around 20% for phenobarbital (Calle et al., 2023; de Groot, 2022; Woodruff and Botto, 2022). Further studies are needed to establish a gold standard method to identify the culprit drug of DRESS syndrome.

Diagnosis of DRESS syndrome

DRESS syndrome is considered to be a challenging diagnosis, mainly due to its heterogeneous clinical features (Calle et al., 2023). The Registry of Severe Cutaneous Adverse Reactions (RegiSCAR) and the Japanese Consensus Group (J-SCAR) are three different diagnostic scoring systems that use distinct sets of criteria. Bocquet et al., (1996) were the first to propose diagnostic criteria for DRESS. These included drug-induced rash, eosinophilia $>1500 \times 10^9/L$, presence of atypical lymphocytes, and systemic involvement such as lymphadenopathy, hepatic, renal, lung, or cardiac manifestations. The RegiSCAR group later suggested another set of diagnostic criteria, in which cases were categorized as „no“, „possible“, „probable“ or „definite“ DRESS (Kardaun et al., 2007). However, this standardized scoring system did not include HHV6 reactivation. Studies have reported that although the RegiSCAR criteria are well-suited for clinical trials, Bocquet's criteria seem more useful in daily practice. Kim and Koh, (2014) proposed that these two scoring systems may be used in a complementary manner. Shiohara et al., (2006) and the Japanese Consensus Group (J-SCAR) established another set of diagnostic criteria. Instead of DRESS, they prefer to use the acronym DIHS. This DIHS scoring system consists of seven criteria: maculopapular rash occurring more than three weeks after starting therapy, persistence of clinical symptoms two weeks after causative drug withdrawal, fever ($>38^\circ C$), liver alterations ($ALT >100/L$), at least one of leukocyte alterations (leukocytosis ($>11 \times 10^9/L$), atypical lymphocytosis ($>5\%$), eosinophilia ($>1.5 \times 10^9/L$)), lymphadenopathy and HHV-6 reactivation. It is important that J-SCAR only includes some drugs as potential triggers for DIHS: carbamazepine, phenytoin, phenobarbital, mexiletine, sulfasalazine, allopurinol and minocycline (Criado et al., 2025). Moreover, compared with the previously mentioned scoring systems, it considers reactivation of viruses in the Herpesviridae family. To establish typical DIHS, all 7 criteria must be met; however, a diagnosis of a modified, atypical version of DIHS can be confirmed by the presence of only the first 5 criteria (Shiohara et al., 2006; López-Rocha et al., 2014). Recent studies compared RegiSCAR and Japanese consensus scoring systems.

Sasidharanpillai et al., (2022) suggested that RegiSCAR criteria should be used for diagnosis of DRESS/DIHS, while reactivation of HHV-6, HHV-7, CMV and EBV should rather be treated as prognostic factors. The standardized literature on DRESS syndrome management is scarce. Brügger et al., (2024) conducted an international expert consensus and reported key aspects of DRESS diagnosis, management, as well as proposed disease severity scale. They established generalized diagnostic workup which should be applicable to all patients suspected with DRESS. It includes complete differential blood count, kidney and liver function tests, urine sediment analysis and electrocardiography. Additionally in patients with suspected specific organ manifestations, more targeted tests should be performed. It can be coagulation studies, lesional skin or liver biopsy, cardiac biomarkers (troponin T, BNP or NT-proBNP), transthoracic echocardiography, pancreatic enzymes, chest or brain imaging, bronchoscopy and lumbar puncture. Brügger et al., (2024) also recommend assessment for HHV-6, EBV, CMV reactivation. HHV-7, hepatitis B virus and hepatitis C virus may be checked in selected groups of patients. There is a paucity of literature on the classification of DRESS syndrome severity. Brügger et al., (2024) suggested the first DRESS severity scale. Cases are categorized into mild, moderate, or severe based on liver, renal, and hematologic markers. Patients with any other organ involvement are classified as severe cases.

How to differentiate DRESS syndrome?

Diagnosis and differentiation of DRESS is challenging due to its high degree of heterogeneity in its clinical presentation. However, patients presenting with classic, typical disease progression, maculopapular rash, fever, and lymphadenopathy that occur weeks after administration of a culprit drug generally do not pose a significant diagnostic challenge (Criado et al., 2025). Conducting the recommended diagnostic workup and taking the patient's history can help with differential diagnosis (Brügger et al., 2024). Stevens-

Johnson syndrome (SJS), toxic epidermal necrolysis (TEN), extensive erythema multiforme, acute generalized exanthematous pustulosis (AGEP), and staphylococcal scalded skin syndrome are the most common diseases that may present with a clinical picture similar to DRESS. Clinicians should also be aware of viral conditions such as EBV infection, CMV, hepatitis A and B, acute HIV infection, COVID-19, and parvovirus B19 (Calle et al., 2023). Dondi et al., (2021) observed cases of DRESS-like viral rash with midfacial edema and systemic symptoms in children treated with amoxicillin during viral illnesses (EBV-infectious mononucleosis (IM) and parainfluenza virus). Moreover, to diagnose DRESS syndrome, specialists must rule out other differentials such as autoimmune and rheumatologic conditions and hematologic malignancies. Clinicians must also distinguish DRESS from other skin-involving diseases, such as vasculitis.

Treatment

The most important step in DRESS management is early discontinuation of potentially causal drugs (Manieri et al., 2023; Choudhary et al., 2013). Clinicians should be aware that DRESS treatment should start as soon as possible, as it can improve the prognosis (de Groot, 2022). Treatment is essentially supportive and systemic. In patients with moderate to severe DRESS, systemic corticosteroids seem to have shown the best outcome in recovery and are considered to be a first-line medication in the acute phase of the disease. However, it should be noted that there are no randomized controlled trials guiding the use of steroids in DRESS, and this therapeutic approach is largely based on clinical opinions.

Nevertheless, they are still widely considered the most optimal treatment method (Calle et al., 2023). In general, it is recommended to start with prednisolone (or equivalent) 0.5–1 mg/kg/day with gradual tapering over 2–3 months to prevent severe or fatal relapses (Shiohara and Kano, 2017). Clinicians should be aware that, besides a slow steroid-tapering regimen, patients with DRESS, in some cases, may need to have them administered for a year or longer due to the increased risk of autoimmune DRESS complications (Ushigome et al., 2013). The results of the study by Santiago et al., (2010) revealed that all 30 patients with DRESS syndrome treated with systemic corticosteroids showed dramatic improvements in symptoms and laboratory results. Moreover, they didn't develop any secondary skin or soft-tissue infections. In mild cases of DRESS, topical steroids and antihistamines are recommended (Hama et al., 2022).

Patients who are steroid-resistant or who cannot tolerate steroids can be treated with different options such as cyclosporine, mycophenolate mofetil, or cyclophosphamide (Wang et al., 2024; Giri et al., 2011). Cyclosporine has been reported as a potential replacement for steroids in immunosuppressive therapy in DRESS patients. However, only a limited number of cases have reported this line of treatment in the literature. Studies have shown that cyclosporine may be successfully used either in patients previously treated with steroids or as a steroid-sparing agent. Typically, doses range from 3 to 5 mg/kg/d. Zita et al., (2023) reported that cyclosporine was also an effective treatment for patients who couldn't tolerate prolonged corticosteroid therapy.

Another drug that may be used in DRESS treatment is Mycophenolate mofetil, an immunosuppressive medication that selectively targets lymphocytes. However, specialists should be aware of its potential side effects, which include gastrointestinal, hematologic, and genitourinary systems (Finelli et al., 2006). Although there is scarce information in the literature about the efficacy of cyclophosphamide in severe DRESS, it is still considered a potential steroid alternative, especially for patients with kidney failure and myocarditis (Wang et al., 2024).

For steroid-resistant and steroid-dependent DRESS patients, intravenous immunoglobulin (IVIG) (Hung et al., 2024; Ushigome et al., 2013), anti-IL-5/anti-IL-5R biologics, and tumor necrosis factor TNF- α are other DRESS treatment options (Wang et al., 2024). Brügggen et al., (2024), following Japanese, Spanish, and French guidelines, agreed that ganciclovir or valganciclovir should be considered in patients with a high CMV viral load or life-threatening CMV-related manifestations.

Follow-up care

Long-term autoimmune sequelae of DRESS syndrome have been reported in the literature. They may be a continuation of the clinical manifestations of the acute phase or may occur after disease remission. Studies have reported that the interval between the acute phase of DRESS and the onset of sequelae can be as long as 4 years. Complications developing within the first weeks after DRESS clinical manifestations include autoimmune thyroiditis, fulminant type 1 diabetes mellitus, fulminant hepatic failure, autoimmune hemolytic anemia, renal failure, and disseminated intravascular coagulation. Arthralgia (including rheumatoid arthritis), vitiligo, alopecia areata, and systemic lupus erythematosus may persist for months to years. Hama et al., (2022) and Brügggen et al., (2024), in an international consensus, suggested that patients should attend follow-up consultations within the first month and be closely monitored during the

first 6 months after DRESS initiation. Close monitoring of patients should include screening for autoantibodies and thyroid dysfunction, screening for steroid adverse effects, as well as active offering of psychological support.

4. CONCLUSION

DRESS syndrome is a rare but severe condition, characterized by rash, fever, systemic and hematological symptoms, and lymphadenopathy. Clinicians should be aware that prompt recognition and discontinuation of the potential medication causing DRESS plays a key role in the patient's treatment. As significant chronic sequelae may occur, long-term monitoring is required. Given the limited literature on optimal management strategies for DRESS, there remains a need for established, standardized guidelines for clinicians.

Abbreviations used:

CMV - Cytomegalovirus

DIHS/DRESS - Drug-induced hypersensitivity syndrome/drug reaction with eosinophilia and systemic symptoms

HHV - Human herpesvirus

J-SCAR - Japanese SCAR

RegiSCAR - Registry of Severe Cutaneous Adverse Reactions to Drugs and Collection of Biological Samples

FAERS- FDA's Adverse Event Reporting System

FDA- Food and Drug Administration

HLA- Human Leukocyte Antigen

SCAR - Severe cutaneous adverse drug reaction

SJS/TEN - Stevens-Johnson syndrome and toxic epidermal necrolysis

CYP- Cytochrome P450

Acknowledgments

We thank all co-authors and institutional staff for their support during the preparation of this review.

Authors' Contributions

Conceptualization and Methodology: Zofia Rogowska, Kaja Pruszkowska, Barbara Olędzka, Małgorzata Olędzka, Aleksandra Rzepko

Check: Sara Omid, Wiktoria Mazepa, Michał Piątkiewicz

Formal analysis: Zofia Rogowska, Aleksandra Rzepko, Małgorzata Olędzka

Investigation: Zofia Rogowska, Sara Omid, Michał Piątkiewicz, Aleksandra Rzepko

Data curation: Zofia Rogowska, Barbara Olędzka, Wiktoria Mazepa, Kaja Pruszkowska

Writing-rough preparation: Zofia Rogowska

Writing review and editing: Sara Omid, Michał Piątkiewicz, Wiktoria Mazepa, Małgorzata Olędzka, Kaja Pruszkowska, Barbara Olędzka, Aleksandra Rzepko

Visualization, Supervision & Project administration: Zofia Rogowska

All authors have reviewed and approved the final version of the manuscript and agree to be accountable for all aspects of the work. All authors contributed equally to the manuscript.

Informed consent

Not applicable.

Ethical approval

Not applicable. This article does not contain any studies with human participants or animals performed by any of the authors.

Funding

This research did not receive any external funding like specific grant from funding agencies in the public, commercial, or nonprofit sectors.

Conflict of interest

The authors declare that they have no conflicts of interest, competing financial interests or personal relationships that could have influenced the work reported in this paper.

Data and materials availability

All data associated with this study will be available based on the reasonable request to corresponding author.

REFERENCES

- Bocquet H, Bagot M, Roujeau JC. Drug-induced pseudolymphoma and drug hypersensitivity syndrome (Drug Rash with Eosinophilia and Systemic Symptoms: DRESS). *Semin Cutan Med Surg* 1996;15(4):250-7. doi: 10.1016/s1085-5629(96)80038-1.
- Brüggen M, Walsh S, Ameri MM. Management of Adult Patients With Drug Reaction With Eosinophilia and Systemic Symptoms: A Delphi-Based International Consensus. *JAMA Dermatol* 2024;160(1):37-44. doi: 10.1001/jamadermatol.2023.4450.
- Cabañas R, Calderón O, Ramírez E, Fiandor A, Caballero T, Heredia R, Herranz P, Madero R, Quirce S, Bellón T. Sensitivity and specificity of the lymphocyte transformation test in drug reaction with eosinophilia and systemic symptoms causality assessment. *Clin Exp Allergy* 2018;48(3):325-333. doi: 10.1111/cea.13076.
- Cacoub P, Musette P, Descamps V, Meyer O, Speirs C, Finzi L, Roujeau JC. The DRESS syndrome: a literature review. *Am J Med* 2011;124(7):588-97. doi: 10.1016/j.amjmed.2011.01.017.
- Calle AM, Aguirre N, Ardila JC, Cardona Villa R. DRESS syndrome: A literature review and treatment algorithm. *World Allergy Organ J* 2023;16(3):100673.
- Choudhary S, McLeod M, Torchia D, Romanelli P. Drug Reaction with Eosinophilia and Systemic Symptoms (DRESS) Syndrome. *J Clin Aesthet Dermatol* 2013;6(6):31-7.
- Criado PR, Ianhez M, Miot HA, Criado RFJ, Talhari C, Müller Ramos P. DRESS syndrome: an interaction between drugs, latent viruses, and the immune system. *An Bras Dermatol* 2025;100(1):104-120. doi: 10.1016/j.abd.2023.12.010.
- de Groot AC. Patch testing in drug reaction with eosinophilia and systemic symptoms (DRESS): a literature review. *Contact Dermatitis* 2022;86(6):443-79. doi: 10.1111/cod.14090.
- Del Pozzo-Magaña BR, Rieder MJ, Garcia-Bournissen F, Lazo-Langner A. Drug reaction with eosinophilia and systemic symptoms (DRESS): A tertiary care centre retrospective study. *Br J Clin Pharmacol* 2022;88(9):4134-4141. doi: 10.1111/bcp.15354.
- Dondi A, Parladori R, Mori F, Liccioli G, Bassi A, Lanari M, Patrizi A, Neri I. Viral rashes mimicking drug reaction with eosinophilia and systemic symptoms syndrome in children after β -lactams intake: a diagnostic challenge. *Eur J Pediatr* 2021;180(7):2327-2332. doi: 10.1007/s00431-021-04010-5.
- Eshki M, Allanore L, Musette P, Milpied B, Grange A, Guillaume JC, Chosidow O, Guillot I, Paradis V, Joly P, Crickx B, Ranger-Rogez S, Descamps V. Twelve-year analysis of severe cases of drug reaction with eosinophilia and systemic symptoms: a cause of unpredictable multiorgan failure. *Arch Dermatol* 2009;145(1):67-72. doi: 10.1001/archderm.145.1.67.
- Finelli PF, Naik K, DiGiuseppe JA, Prasad A. Primary lymphoma of CNS, mycophenolate mofetil and lupus. *Lupus* 2006;15(12):886-8. doi: 10.1177/0961203306071431.
- Giri PP, Roy S, Bhattiyacharya S, Pal P, Dhar S. Dress syndrome with sepsis, acute respiratory distress syndrome and pneumomediastinum. *Indian J Dermatol* 2011;56(6):763-5. doi: 10.4103/0019-5154.91850.
- Hama N, Abe R, Gibson A, Phillips EJ. Drug-Induced Hypersensitivity Syndrome (DIHS)/Drug Reaction With Eosinophilia and Systemic Symptoms (DRESS): Clinical Features and Pathogenesis. *J Allergy Clin Immunol Pract* 2022;10(5):1155-67.e5. doi: 10.1016/j.jaip.2022.02.004.
- Hiransuthikul A, Rattananupong T, Klaewsongkram J, Rerknimitr P, Pongprutthipan M, Ruxrungtham K. Drug-induced hypersensitivity syndrome/drug reaction with eosinophilia and systemic symptoms (DIHS/DRESS): 11 years retrospective study in Thailand. *Allergol Int* 2016;65(4):432-8. doi: 10.1016/j.alit.2016.04.001.
- Hung SI, Chung WH, Liou LB, Chu CC, Lin M, Huang HP, Lin YL, Lan JL, Yang LC, Hong HS, Chen MJ, Lai PC, Wu MS, Chu CY, Wang KH, Chen CH, Fann CS, Wu JY, Chen YT. HLA-B*5801 allele as a genetic marker for severe cutaneous adverse reactions caused by allopurinol. *Proc Natl Acad Sci U S A* 2005;102(11):4134-9. doi: 10.1073/pnas.0409500102.
- Hung SI, Mockenhaupt M, Blumenthal KG, Abe R, Ueta M, Ingen-Housz-Oro S, Phillips EJ, Chung WH. Severe cutaneous adverse reactions. *Nat Rev Dis Primers* 2024;10(1):30. doi: 10.1038/s41572-024-00514-0.
- Kardaun SH, Sekula P, Valeyrie-Allanore L, Liss Y, Chu CY, Creamer D, Sidoroff A, Naldi L, Mockenhaupt M, Roujeau JC;

- RegiSCAR study group. Drug reaction with eosinophilia and systemic symptoms (DRESS): an original multisystem adverse drug reaction. Results from the prospective RegiSCAR study. *Br J Dermatol* 2013;169(5):1071-80. doi: 10.1111/bjd.12501.
19. Kardaun SH, Sidoroff A, Valeyrie-Allanore L, Halevy S, Davidovici BB, Mockenhaupt M, Roujeau JC. Variability in the clinical pattern of cutaneous side-effects of drugs with systemic symptoms: does a DRESS syndrome really exist? *Br J Dermatol* 2007;156(3):609-11. doi: 10.1111/j.1365-2133.2006.07704.x.
 20. Kim DH, Koh YI. Comparison of diagnostic criteria and determination of prognostic factors for drug reaction with eosinophilia and systemic symptoms syndrome. *Allergy Asthma Immunol Res* 2014;6(3):216-21. doi: 10.4168/aaair.2014.6.3.216.
 21. Kroshinsky D, Cardones ARG, Blumenthal KG. Drug Reaction with Eosinophilia and Systemic Symptoms. *N Engl J Med* 2024;391(23):2242-2254. doi: 10.1056/NEJMra2204547.
 22. Lehloenya RJ, Peter JG, Copascu A, Trubiano JA, Phillips EJ. Delabeling delayed drug hypersensitivity: how far can you safely go? *J Allergy Clin Immunol Pract* 2020;8(9):2878-95.e6. doi: 10.1016/j.jaip.2020.07.005.
 23. Liang C, An P, Zhang Y, Liu X, Zhang B. Fatal outcome related to drug reaction with eosinophilia and systemic symptoms: a disproportionality analysis of FAERS database and a systematic review of cases. *Front Immunol* 2024;15:1490334. doi: 10.3389/fimmu.2024.1490334.
 24. López-Rocha E, Blancas L, Rodríguez-Mireles K, Gaspar-López A, O'Farrill-Romanillos P, Amaya-Mejía A, Galindo-Pacheco L, Campos-Romero F, Aguilar-Hinojosa N, Suárez G. Prevalence of DRESS syndrome. *Rev Alerg Mex* 2014;61(1):14-23.
 25. Manieri E, Dondi A, Neri I, Lanari M. Drug rash with eosinophilia and systemic symptoms (DRESS) syndrome in childhood: a narrative review. *Front Med (Lausanne)* 2023;10:1108345. doi: 10.3389/fmed.2023.1108345.
 26. Miyagawa F, Asada H. Current Perspective Regarding the Immunopathogenesis of Drug-Induced Hypersensitivity Syndrome/Drug Reaction with Eosinophilia and Systemic Symptoms (DIHS/DRESS). *Int J Mol Sci* 2021;22(4):2147. doi: 10.3390/ijms22042147.
 27. Peter JG, Lehloenya R, Dlamini S, Risma K, White KD, Konvinse KC, Phillips EJ. Severe Delayed Cutaneous and Systemic Reactions to Drugs: A Global Perspective on the Science and Art of Current Practice. *J Allergy Clin Immunol Pract* 2017;5(3):547-63. doi: 10.1016/j.jaip.2017.01.025.
 28. Rahnema-Moghadam S, Arora N, Vuppalachchi R, Li YJ, Gu J, Barnhart H, Phillips E, Chalasani N; Drug Induced Liver Injury Network. DRESS Syndrome in Patients With Drug-Induced Liver Injury: Characteristics and HLA Risk Factors. *Am J Gastroenterol* 2025. doi: 10.14309/ajg.0000000000003590.
 29. Santiago F, Gonçalves M, Vieira R, Coelho S, Figueiredo A. Epicutaneous patch testing in drug hypersensitivity syndrome (DRESS). *Contact Dermatitis* 2010;62(1):47-53. doi:10.1111/j.1600-0536.2009.01659.x.
 30. Sasidharanpillai S, Ajithkumar K, Jishna P, Khader A, Anagha KV, Binitha MP, Chathoth AT. RegiSCAR DRESS (Drug Reaction with Eosinophilia and Systemic Symptoms) Validation Scoring System and Japanese Consensus Group Criteria for Atypical Drug-Induced Hypersensitivity Syndrome (DIHS): A Comparative Analysis. *Indian Dermatol Online J* 2022;13(1):40-45. doi: 10.4103/idoj.idoj_196_21.
 31. Shiohara T, Inaoka M, Kano Y. Drug-induced hypersensitivity syndrome (DIHS): A reaction induced by a complex interplay among herpesviruses and antiviral and antidrug immune responses. *Allergol Int* 2006;55(1):1-8. doi: 10.2332/allergolint.55.1.
 32. Shiohara T, Kano Y. Drug reaction with eosinophilia and systemic symptoms (DRESS): incidence, pathogenesis and management. *Expert Opin Drug Saf* 2017;16(2):139-147. doi: 10.1080/14740338.2017.1270940.
 33. Shiohara T, Mizukawa Y. Drug-induced hypersensitivity syndrome (DIHS)/drug reaction with eosinophilia and systemic symptoms (DRESS): An update in 2019. *Allergol Int* 2019;68(3):301-8. doi: 10.1016/j.alit.2019.03.006.
 34. Stirton H, Shear NH, Dodiuk-Gad RP. Drug Reaction with Eosinophilia and Systemic Symptoms (DRESS)/Drug-Induced Hypersensitivity Syndrome (DIHS)-Readdressing the DRESS. *Biomedicine* 2022;10(5):999. doi: 10.3390/biomedicine10050999.
 35. Taweessdt PT, Nordstrom CW, Stoeckel J, Dumić I. Pulmonary Manifestations of Drug Reaction with Eosinophilia and Systemic Symptoms (DRESS) Syndrome: A Systematic Review. *Biomed Res Int* 2019;2019:7863815. doi: 10.1155/2019/7863815.
 36. Toniato A, Gamba C, Schroeder JW, Fabbri V, Bernal Ortiz SV, Borgonovo L, Piantanida M, Scibilia J, Balossi L, Brusamolino E, Bonoldi E, Caputo V, Nichelatti M, Pastorello EA. Role of Multiple Comorbidities and Therapies in Conditioning the Clinical Severity of DRESS: A Mono-Center Retrospective Study of 25 Cases. *Int J Mol Sci* 2021;22(13):7072. doi: 10.3390/ijms22137072.
 37. Ushigome Y, Kano Y, Ishida T, Hirahara K, Shiohara T. Short- and long-term outcomes of 34 patients with drug-induced hypersensitivity syndrome in a single institution. *J Am Acad Dermatol* 2013;68(5):721-8. doi: 10.1016/j.jaad.2012.10.017.

38. Wang L, Mei XL. Drug reaction with eosinophilia and systemic symptoms: retrospective analysis of 104 cases over one decade. *Chin Med J (Engl)* 2017;130(8):943-9. doi:10.4103/0366-6999.204104.
39. Wang S, Kang Y, He C, Jin H. The systemic treatments for drug reaction with eosinophilia and systemic symptoms (DRESS) beyond corticosteroids. *World Allergy Organ J* 2024; 17(8):100935. doi: 10.1016/j.waojou.2024.100935.
40. Woodruff CM, Botto N. The role of patch testing in evaluating delayed hypersensitivity reactions to medications. *Clin Rev Allergy Immunol* 2022;62(3):548-61. doi: 10.1007/s12016-022-08924-2.
41. Zita S, Broussard L, Hugh J, Newman S. Cyclosporine in the Treatment of Drug Reaction with Eosinophilia and Systemic Symptoms Syndrome: Retrospective Cohort Study. *JMIR Dermatol* 2023;6:e41391. doi: 10.2196/41391.