

## SPECIES

## To Cite:

Ghory FS, Tahera Q, Ríos-Escalante PRDL. Morphotaxonomy of shallow water *Comanthus wahlbergii* (Müller, 1843) (Echinodermata: Crinoidea: Comasteridae) collected from Jiwani (Makran coast, Pakistan). *Species* 2023; 24: e51s1545  
doi: <https://doi.org/10.54905/disssi/v24i73/e51s1545>

## Author Affiliation:

<sup>1</sup>Marine Reference Collection and Resource Centre, University of Karachi, Karachi-75270, Pakistan

<sup>2</sup>Departamento de Ciencias Biológicas y Químicas, Facultad de Recursos Naturales, Universidad Católica de Temuco, Casilla (PO-Box) 15-D, Temuco, Chile

<sup>3</sup>Núcleo de Estudios Ambientales, Universidad Católica de Temuco, Temuco, Chile

## Corresponding author

Marine Reference Collection and Resource Centre, University of Karachi, Karachi-75270, Pakistan  
Email: [farhanaghory@yahoo.com](mailto:farhanaghory@yahoo.com)

## Peer-Review History

Received: 09 April 2023

Reviewed & Revised: 14/April/2023 to 05/June/2023

Accepted: 08 June 2023

Published: 15 June 2023

## Peer-Review Model

External peer-review was done through double-blind method.

## Species

pISSN 2319-5746; eISSN 2319-5754



© The Author(s) 2023. Open Access. This article is licensed under a [Creative Commons Attribution License 4.0 \(CC BY 4.0\)](https://creativecommons.org/licenses/by/4.0/), which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons license, and indicate if changes were made. To view a copy of this license, visit <http://creativecommons.org/licenses/by/4.0/>



# Morphotaxonomy of shallow water *Comanthus wahlbergii* (Müller, 1843) (Echinodermata: Crinoidea: Comasteridae) collected from Jiwani (Makran coast, Pakistan)

Farhana S Ghory<sup>1\*</sup>, Qaseem Tahera<sup>1</sup>, Patricio R De Los Ríos-Escalante<sup>2,3</sup>

## ABSTRACT

Crinoidea is the class of marine animals composed of crinoids, which comprises the feather-stars or sea-lilies. Vast arrays of living and extinct species are found in shallow water to great depths in this most primitive of living classes of Echinodermata (Pandian, 2018). Many species of stalked crinoids live in deep seas and a few are sessile, which are exhibited on rocky beaches. There are a variety of colors, from the most vibrant animals to the pale or colorless. Only the genus *Comanthus* A.H.Clark, 1908 is found in Pakistani waters. Presently we described the taxonomy of shallow-water *Comanthus wahlbergii* (Müller, 1843) that were collected from Jiwani (Pakistan) (25°03'06"N 61°44'35"E).

**Keywords:** *Comanthus wahlbergii*, shallow water, feather stars, Jiwani (Makran coast, Pakistan).

## 1. INTRODUCTION

In the marine environment, echinoderms are among the most widespread and common animal groups (Hyman, 1955; Hickman et al., 2023). Additionally, they are clearly defined among animal phyla. The members of this group have basic pentamerous adults, a calcite skeleton and hydraulic tube feet connected to a water vascular system (Rowe and Gates, 1995; Hickman et al., 2023). There has been significant evolution within the phylum of echinoderms due to the unique characteristics of their skeletal systems (Lawrence, 1987; Hickman et al., 2023).

Echinoderms comprise animals usually known as feather star and sea lilies (Crinoidea); starfish (Asteroidea); brittle stars (Ophiuroidea); sea urchins, sand dollars and heart urchins (Echinoidea); and sea cucumbers (Holothuroidea). There are several commercial marine animals that work in marine ecosystems and have significant role in the overall ecology of the ocean as sediment recyclers,

scavengers, burrowers and predators (Hyman, 1955; Rowe and Gates, 1995; Putchakarn, 1998; Hickman et al., 2023). There are a variety of microhabitats for symbiotic fauna on crinoids, particularly polychaetes, myzostomids, shrimps and crabs, due to their complex external morphology (Sorokin, 1990; Fabricius and Dale, 1993; Deheyn et al., 2006; Britayev and Mekhova, 2011; Hickman et al., 2019).

Crinoids are considered to be a keystone group in tropical coral reefs because of their many symbiotic associations. Biodiversity studies of marine ecosystems tend to overlook and underestimate this symbiotic fauna. Crinoids filter plankton and tiny detritus particles out of flowing sea water with their feather-like arms. The arms are raised and confined vertical to the current to form a fan shape. The identification of crinoids is a difficult task. Patience and attention to skeletal details are necessary. The present study investigated the taxonomy of *Comanthus wahlbergii* collected from shallow water of Makran coast (Jiwani).

## 2. MATERIALS AND METHODS

Sixteen specimens were obtained all through low tide from intertidal zone of Jiwani (Makran coast, Pakistan) (Lat. 25°03'06''N Long. 61°44'35''E). For observation of crinoids were relaxed by spreading them in sea water, aqueous MgSO<sub>4</sub> (Epsom salt) was found to be the best narcotizing agent. The specimens were then spread out in a flat pan or a large dish with sea water, their arms not touching one another, MgSO<sub>4</sub> was sprinkled and fresh water was added gradually. The fully relaxed specimens were finally preserved in 70% alcohol. Length of arms 130 - 151mm. Radius of centrodorsal 4.0 - 6.0mm. Systematic was reviewed in according to WoRMS, (2023).

### Systematic

Phylum Echinodermata Bruguière, 1791

Subphylum Crinozoa Matsumoto 1929

Class Crinoidea Miller, 1821

Subclass Articulata Zittel, 1879

Order Comatulida A. H. Clark, 1908

Superfamily Comatuloidea Fleming, 1828

Family Comatulidae Fleming, 1928

Subfamily Comatulinae Fleming, 1828

Genus *Comanthus* A.H. Clark, 1908

Species *Comanthus wahlbergii* (Müller, 1843)

***Comanthus wahlbergii* (Müller, 1843) (Figure 1, 2)**

*Comanthus wahlbergii* (Müller, 1843)

*Alecto wahlbergii* (Müller, 1843).

*Comanthus samoana* (Clark, 1912; Clark, 1913).

*Comanthus* (*Comanthus*) *samoana* (Clark, 1909).

*Comanthus samoanus* (Clark, 1909; Clark and Rowe, 1971).

*Comanthus wahlbergi serratus* (Clark, 1972).

*Comanthus wahlbergi tenuibrachia* (Clark, 1972) (Figure 1).

*Comissia serrata* (Gislén, 1938).

*Comanthus walbergii* (Clark, 1931; Clark, 1972; Rowe et al., 1986).



**Figure 1** *Comanthus wahlbergii* (Müller, 1843)

### 3. RESULTS AND DESCRIPTION

The species has previously been reported from Jiwani by Hoque, (1969) as = *Comanthus samoanus*. The present specimens afford the first consequent record from this region. *Comanthus wahlbergii* (Figure 2A) is characterized by having 21 arms measuring 45.0mm in length and only 3.0mm diameter at the first syzygy. The centrodorsal is long compressed and discoidal it is 6.0mm in diameter and with one inclusive marginal row of XXIV-XXVI cirri.

All cirrus (Figure 2B) is stout, curved, jointed and well developed; arising from the centrodorsal measuring about 16.0mm in length and having up to 14 segments. The first 3 or 4 segments are small, the last 5 or 6 with dorsal crest. The penultimate segment bears stout, curved terminal claw. The brachial and pinnules are densely spinose. The proximal pinnules (Figure 2C) are exceedingly flexible with some of the terminal segments modified to form a comb. The combs occur at intervals extending up to the tips of the arms. The mouth is close to the edge of the disc. The anal tube is more or less central.

#### Color


Live specimens with dark brown aboral surface. Oral surface and cirri pale brown.

#### Ecology

An abundant, diurnal species found on boulders, incline surfaces and hard corals. A cirri is used to fix them to the surface. The arms of the filtration fan form a biplanar shape during feeding. Arrangement of pinnules in four planes. The tips of the inactive arms were pointing towards the calyx. Like environments with strong currents. Found at 3–5 m.

#### Distribution

Pacific Ocean, Pakistan, India, Sri Lanka, Red Sea, East Africa, South Africa, Madagascar, Seychelles, South-China Sea, Vietnam, Japan, north Australia, Great Barrier Reef, New Caledonia, New Zealand and Philippine.



**Figure 2** *Comanthus wahlbergii* (Müller, 1843). A) Entire specimen, dorsal view; B) Single cirrus; C) Single first pinnule

#### 4. CONCLUDING REMARKS

There are more than 150 shallow-water species of unstalked crinoids in the Indian Ocean and western Pacific Ocean, with the greatest diversity (Messing, 1997). They may be able to avoid predators and find food since most of these organisms are not fixed to the substrate (Meyer and Macurda, 1977). Studies of crinoids' systematic, phylogenies and fauna have been conducted in great detail (Clark and Rowe, 1971; Rowe et al., 1986; Messing and White, 2001; Messing, 2003; Cohen et al., 2004). In spite of this, the fauna of a variety of areas of the World Ocean is poorly understood.

In particular, the fauna of the Palau Islands (Meyer and Macurda, 1980), Great Barrier Reef of Australia (Messing, 1994), Japan (Kogo, 1998; Kogo, 2006) and the Marshall Islands (Zmarzly, 1984; Zmarzly, 1985) are quite wholly investigated. Family comasteridae contains the subsequent eight genera reported from Indo-West Pacific region: *Capillaster*, *Comantheria*, *Comanthina*, *Comanthus*, *Comaster*, *Comatella*, *Comatula* and *Comissia* (Clark and Rowe, 1971). Only one genus *Comanthus* is found in Pakistan. There are only two recognized species of the genus *Comanthus*: *C. wahlbergii* (Müller, 1843) and *C. parvicirrus* Müller, 1841. Only *C. wahlbergii* is represented in our collection.

This species exhibits many of the adaptations and behaviors that have allowed it to exploit shallow marine habitats around the globe (Meyer and Macurda, 1977). They are capable to re-grow mainly of their organs after loss or damage, as well as arms and viscerae (Mozzi et al., 2006) and they have numerous morphological traits that minimize damage during predation trial (Oji and Okamoto, 1994). In

particular, feather stars, as well as *Comanthus wahlbergii*, typically branch the majority at the bottom of the rays (Oji and Okamoto, 1994). According to Baumiller and Gahn, (2013), this species loses an arm every 10 days. We lack long term observations of the crinoid populations. It is expected that further studies will expand the variety of crinoid species known in Pakistan's waters.

**Table 1** Morphological differences between *C. wahlbergii* (Müller, 1843) and *C. parvicirrus* J.Müller, 1841

S. No.	<i>C. wahlbergii</i> (Müller, 1843)	<i>C. parvicirrus</i> J.Müller, 1841
1	Cirri are well developed, numerous (35–40) arranged in two rows on the edge of the Centro dorsal	Cirri very reduced in size and number, often infrequently set around the edge of the centrodorsal, distal segments only slightly shorter than the proximal ones
2	Less than 40 arms, usually about 20 or 21	Usually 19 – 42 arms
3	IIBr four, rarely two and IIIBr series four	IIBr and IIIBr series usually two

#### Disclosure statement

There is no financial support for this research work from the funding agency.

#### Informed consent

Not applicable.

#### Ethical approval & declaration

In this article, as per the marine animal regulations in Marine Reference Collection and Resource Centre, University of Karachi, Pakistan, the authors observed the morphotaxonomy of shallow water *Comanthus wahlbergii* (Müller, 1843) (Echinodermata: Crinoidea: Comasteridae) from Jiwani (Makran coast, Pakistan). The Animal ethical guidelines are followed in the study for species observation, identification & experimentation.

#### Conflicts of interests

The authors declare that there are no conflicts of interests.

#### Funding

The study has not received any external funding.

#### Data and materials availability

All data associated with this study are present in the paper.

## REFERENCES AND NOTES

- Baumiller TK, Gahn FJ. Reconstructing predation pressure on crinoids: Estimating arm- loss rates from regenerating arms. *Paleobiology* 2013; 39(1):40-51.
- Britayev TA, Mekhova ES. Assessment of hidden diversity of crinoids and their symbionts in the Bay of Nhatrang, Vietnam. *Org Divers Evol* 2011; 11(4):275-285.
- Clark AH. A monograph of the existing crinoids. The Comatulids. Superfamily Comasterida. *Bull U S Natl Mus* 1931; 1(3):82:i-vii, 1-816, pls 1-82.
- Clark AH. Five new species of recent unstalked Crinoids. *Proc U S Natl Mus* 1909; 37:29-34.
- Clark AH. Noted on the recent crinoids in the British Museum. *Smithsonian Miscellaneous Collections* 1913; 61:1-89.
- Clark AH. On a collection of recent crinoids from the Philippine Islands. *Proc U S Natl Mus* 1909; 36:410.
- Clark AH. The crinoids of the Indian Ocean. The Trustees of the Indian Museum, Calcutta 1912; 325, 61 figs.
- Clark AM, Rowe FWE. Monograph of Shallow Water Indo-West Pacific Echinoderms. *Br Mus Nat Hist London* 1971; 238.
- Clark AM. Some crinoids from the Indian Ocean. *Bull Br Mus Nat Hist Zool* 1972; 24:75-156.
- Cohen BL, Ameziane N, Eleaume M, Forges BR. Crinoid Phylogeny: A Preliminary Analysis (Echinodermata: Crinoidea). *Mar Biol* 2004; 144:605-617.
- Deheyn D, Lyskin SA, Eeckhaut I. Assemblages of Symbionts in Tropical Shallow\_Water Crinoids and Assessment of Symbionts' Host\_Specificity. *Symbiosis* 2006; 42:161-168.
- Fabricius KE, Dale MB. Multispecies associations of symbionts on shallow water crinoids of the central Great Barrier Reef. *Coenoses* 1993; 8:41-52.



13. Gislén T. Crinoids of S. Africa. Kungl Vetensk Acad Handl Ser 1938; 3(17):1–22, pls 1–2.
14. Hickman C Jr, Keen CS, Eisenhour D, Larson A, I'Anson H, Cleveland P, Frances M, Lee B. Integrated principles of Zoology (Edition 19). New York: McGraw-Hill 2023.
15. Hickman C Jr, Roberts L, Larson A, I'Anson H. Laboratory Studies in Integrated Principles of Zoology 18th Edition. McGraw Hill 2019; 448.
16. Hoque MM. Echinoderms of Pakistan Coasts. Rec Zool Surv Pak 1969; 1(2):27–38.
17. Hyman LH. The invertebrates. Echinodermata. McGraw- Hill, New York 1955; 4:763.
18. Kogo I. Comatulid fauna (Echinodermata: Crinoidea) of the Sagami Sea and a part of Izu Islands, central Japan. Memoirs of the National Science Museum, Tokyo 2006; 41:223–246.
19. Kogo I. Crinoids from Japan and its adjacent waters. Osaka Museum of Natural History, Special Publication 1998; 30:1–148.
20. Lawrence JMA. Functional Biology of Echinoderms. Johns Hopkins University Press 1987; 340.
21. Messing CG, White CM. A Revision of the Zenometridae (New Rank) (Echinodermata: Crinoidea: Comatulidina). Zool Scr 2001; 30(3):159–180.
22. Messing CG. Comatulid crinoids (Echinodermata) of Madang, Papua New Guinea and environs: Diversity and ecology. In: David B, Guille A, Féral JP, Roux M (Editors): Echinoderms through Time. Proceedings of the Eighth International Echinoderm Conference 1993. Rotterdam 1994; 237–243.
23. Messing CG. Living comatulids. In: Waters JA, Maples CG (Editors) Geology of Echinoderms. The Paleontological Society Papers. Carnegie Museum of Natural History, Pittsburgh 1997; 3:3–30.
24. Messing CG. Three New Species of Comasteridae (Echinodermata: Crinoidea) from the Tropical Western Pacific, Zoosystm 2003; 25(1):149–162.
25. Meyer DL, Macurda DB Jr. Adaptive radiation of the comatulid crinoids. Paleobiology 1977; 3(1):74–82.
26. Meyer DL, Macurda DB Jr. Ecology and distribution of the shallow-water crinoids of Palau and Guam. Micronesica 1980; 16:59–99.
27. Mozzi D, Dolmatov IY, Bonasoro F, Candia-Carnevali MD. Visceral regeneration in the crinoid *Antedon mediterranea*: Basic mechanisms, tissues and cells involved in gut regrowth. Cent Eur J Biol 2006; 1(4):609–635.
28. Müller J. Neue Beiträge zur Kenntnis der Arten der Comatulen. Arch Naturgesch 1843; 9:131–136.
29. Müller J. Über die Gattungen und Arten der Comatulen. Arch Naturgesch 1841; 7:139–148.
30. Oji T, Okamoto T. Arm autonomy and arm branching pattern as anti-predatory adaptations in stalked and stalkless crinoids. Paleobiology 1994; 20(1):27–39.
31. Pandian TJ. Reproduction and Development in Echinodermata and Prochordata. Routledge 2018; 286.
32. Putchakarn S. Taxonomic study on echinoderms from the eastern coast of Thailand. Report submitted to the National Research Council Thailand 1998; 109, 8 pls.
33. Rowe FEW, Gates J. Zoological Catalogue of Australia, Echinodermata. CSIRO, Melbourne 1995; 33:510.
34. Rowe FWE, Hoggett AK, Birtles RA, Vail LL. Revision of some comasterid genera from Australia (Echinodermata: Crinoidea), with descriptions of two new genera and nine new species. Zool J Linn Soc 1986; 86(3):197–277.
35. Sorokin IY. Ekosistemy korallovykh rifov (Ecosystems of Coral Reefs), Moscow: Nauka 1990.
36. WoRMS Editorial Board. World Register of Marine Species 2023. doi: 10.14284/170
37. Zmarzly DL. Distribution and ecology of shallow-water crinoids at Enewetak Atoll, Marshall Island, with an annotated checklist of their symbionts. Pac Sci 1984; 38(2):105–122.
38. Zmarzly DM. The shallow-water crinoid fauna of Kwajalein Atoll, Marshall Islands: Ecological observations, interatoll comparisons and zoogeographic affinities. Pac Sci 1985; 39:340–358.