

Climate Change

To Cite:

Mesike CS, Idoko SO. An overview of Carbon sequestration potential of Rubber tree plantations. *Climate Change* 2024; 10: e10cc1043
doi: <https://doi.org/10.54905/dissci.v10i28.e10cc1043>

Author Affiliation:

Rubber Research Institute of Nigeria, PMB 1049 Benin City, Edo State, Nigeria

*Corresponding Author

Rubber Research Institute of Nigeria, PMB 1049 Benin City, Edo State, Nigeria
Email: sammesike@gmail.com

Peer-Review History

Received: 07 June 2024
Reviewed & Revised: 11/June/2024 to 02/September/2024
Accepted: 05 September 2024
Published: 12 September 2024

Peer-Review Model

External peer-review was done through double-blind method.

Climate Change

pISSN 2394-8558; eISSN 2394-8566

URL: http://www.discoveryjournals.org/climate_change



© The Author(s) 2024. Open Access. This article is licensed under a [Creative Commons Attribution License 4.0 \(CC BY 4.0\)](https://creativecommons.org/licenses/by/4.0/), which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons license, and indicate if changes were made. To view a copy of this license, visit <http://creativecommons.org/licenses/by/4.0/>.

An overview of Carbon sequestration potential of Rubber tree plantations

Chinye S Mesike*, Suleman O Idoko

ABSTRACT

Rubber trees capture atmospheric CO₂ through photosynthesis, reduces Carbon (C) from the atmosphere and store it for several decades in the plant tissues as biomass. The central carbon pools in rubber plantation are biomass that are above ground, below ground and dead organic material. Sequestering atmospheric CO₂ into long-lived wood biomass through afforestation and reforestation is a vital tool to mitigate global warming and climate change. We reviewed the existing information of carbon stocks in rubber tree plantations considering the biomass above ground, biomass below ground, dead organic materials and latex harvested. Past studies on contributions of rubber plantations to climate change mitigation with focus on Carbon stocked in tree biomass above and below ground showed that rubber plantations constitute C stocks comparable to some agroforestry or forestry systems. However, the global Greenhouse Gas (GHG) emissions balance needs to consider the establishment of rubber plantations on previous land use.

Keywords: Carbon sequestration, carbon stock, rubber plantations, land uses, climate change mitigation

1. INTRODUCTION

Rubber (*Hevea brasiliensis* Müll. Arg.) is one of the most economically important tree crops in tropical areas of the world. Rubber plantations provide natural rubber latex and rubber wood for several downstream industries. In addition to giving latex and wood, rubber plantations also help to purify the air we breathe by capturing harmful Carbon Dioxide (CO₂) and reducing Global Warming. Rubber trees grows up to 40 meters (m) high in natural forest but does not exceed 25m in height when cultivated. Rubber tree is a perennial crop by nature that is usually cut-down and replanted after 35 years when latex yields decreases to an uneconomic level (Mesike et al., 2010). According to Warren-Thomas et al., (2018) and Ziegler et al., (2009), there has been significant commercial expansion of natural rubber in the tropical rainforest region. The Southeast Asia countries such as Thailand, Indonesia, Vietnam, and Malaysia produces almost 83% of natural rubber in the world (Blagodatsky et al., 2016).

The expansion in the cultivation of natural rubber is driven by the increase for natural rubber demand in China and India. According to the land area planted for rubber has increased to 14.1 million ha over the last three decades. The increase in the land area planted for natural rubber is visible in the Mekong region and Côte d'Ivoire in Africa. This expansion of natural rubber is expected to increase with a projection of 2.4% per annum in the next decade (IRSG, 2019). Climate change is a major environmental problem in the world. Climate change is defined as the emission of greenhouse gases (CO₂ CH₄ N₂O) as a result of direct or indirect burning of non-renewable resources. According to FAO, (2002), the tropical rainforest contains the most significant living biomass on very delicate soils and it may lose its fertility completely when precise cutting are performed.

According to IPCC, (2006) about 1.5 billion tons of carbon is released into the atmosphere each year due to tropical deforestation for agricultural expansion when forests and grasslands are cleared, burned and converted to agricultural systems. Agriculture contributes to over 20 percent of worldwide anthropogenic greenhouse gas emissions. According to Houghton et al., (2012), land-use system and land-cover change contributed about 33% of the global carbon emissions for the past 150 years. However, the current relative contribution of global carbon emissions has declined to about 13 % annually (Houghton et al., 2012). Notwithstanding, the IPCC has posited that tropical deforestation is estimated to release about 1.9 billion tons of carbon into the atmosphere each year.

There is a growing interest of researcher in the world in reducing the emissions rate of greenhouse gases from agriculture related activities. However, increased land-use system such as forestry and agroforestry systems will significantly reduce the atmospheric greenhouse gas levels for more extended periods. According to Soto-Pinto et al., (2010) and Verchet et al., (2007), tree crop plantations can sequester carbon for more extended periods with minimum annual fluctuations and also have a high significant sequestration potential when compared to regular tropical agriculture. However, Liguori et al., (2009) has argued that annual crop like maize can fix more carbon than tree crops in any given year.

Still, their biomass decomposes very fast and carbon is sequestered to the atmosphere very fast. Although, establishment of rubber plantations can drive deforestation but they have large potentials to sequester atmospheric carbon into the biomass and soil at the end of rotational cycle than other tropical crops such as oil palm, cocoa, citrus (Kongsager et al., 2013; Sun et al., 2017). In addition, rubber production is likely to grow the economies of the tropical countries with relatively low net carbon emissions. The underlying goals of this paper review the information pertaining to biomass carbon, soil organic carbon (SOC) present in rubber tree plantation and the effects of land –use change.

2. CARBON STOCK IN RUBBER PLANTATIONS

The central carbon stocks in rubber plantations are the aboveground biomass (AGB), belowground biomass (BGB), soil organic carbon (SOC), and other minor components such as litter layer and dead wood. Collected latex is also considered when calculating the carbon sequestration potential of rubber plantations. The carbon stocks estimate in rubber plantations can be determined by three main factors: viz, (1) plantation age; (2) plantation management (tree density, latex tapping and fertilization) and (3) environmental and edaphic conditions controlled by the local climate and topography. The third factor is the primary determinant for the size of SOC stocks – an essential part of total ecosystem carbon (TEC). SOC has the most extended lifetime if undisturbed, and it contains historical land use and local edaphic conditions. Carbon sequestration potentials of rubber trees were estimated in several studies over the years.

There are wide variations in the growth of rubber in various tropical and subtropical regions due to different climate patterns, altitudes, and soil types (Brahma et al., 2018; Choudhary et al., 2016; Grieco, 2011; Kongsager et al., 2013; Liu et al., 2017; Maggiotto et al., 2014; Yang et al., 2016). In a separate study that was conducted in China by Cheng et al., (2007) and Song and Zhang, (2010), they found a high carbon sequestration potential in rubber plantations in China. In fact, Cheng et al., (2007) concluded that rubber plantations could sequester about 272.08 tC/ha within a 30-year-life span, while Song and Zhang, (2010) estimated about 123.49 tC/ha carbon stock for a plantation with a higher elevation of about 550-600m. In another experiment conducted in West Africa region, Kongsager et al., (2013) has estimated the biomass in four tree crops in Ghana with focus on rubber, cocoa, oil palm and orange.

The most considerable carbon content of 214 tC/ha was found in rubber plantations while cocoa, orange and oil palm plantations have a much lower carbon content of 65 tC/ha, 76 tC/ha and 45 tC/ha respectively. In an experiment carried out by Day, (2005) to estimate the average carbon stock in 51,510 ha of rubber plantations in North-Eastern states of India, he observed that the average carbon stored in rubber plantations was about 136tC/ha out of which 92.7tC/ha was from soil and 2.40tC/ha was through litter fall and undergrowth vegetation. In another work carried out in Xishuangbanna, China by Nizami et al., (2014), to study the effects of rotation

length on the C stocks in rubber plantations, he found that the total carbon stocks was 186.65 Mg C/ha and concluded that more extended rotational lengths have higher carbon stocks in below-ground carbon stock after four consecutive rotations.

In a similar experiment by Yang et al., (2017) and Yang et al., (2019), about 148Mg C/ha carbon stock was estimated in old rubber plantations in Xishuangbanna, China. In the conclusion of the study conducted by Kiyono et al., (2014), in Northern Laos, it was posited that carbon stock of 50.0Mg C/ha in the biomass of a rubber trees of economic life of 30 years was much greater than carbon stock of 18.6Mg C/ha of a five years fallow period slash-and-burn agricultural system.

3. EFFECT OF LAND USE CHANGE

Song and Zhang, (2010) have argued that land use by cultivating rubber plantations in the short-term leads to carbon sequestration. Also, Fox et al., (2014) have also argued that land use change from swiddening (slash and burn) agriculture to a rubber plantation can also lead to carbon but, carbon stocks in land use types vary and dependent on the development stage, management type, location etc of the plantation (Ziegler et al., 2012). According to Fox et al., (2014), Rotational swiddening agriculture were classified into fallow phases of different lengths, such as short length of less than five years; medium length of 5 to 10 years and long length of 10 to 25 years. Depending on the type of length, the amount of carbon sequestration for a land use type can vary. According to Petsri et al., (2013), conversion of forest to rubber plantation usually results to strong loss of C.

In another study carried out by Ziegler et al., (2012), they observed that there was a slight mean positive trend for long-term land fallow when swiddening agriculture is substituted with rubber plantation. They concluded that when a secondary forest is converted to a rubber plantation for prolonged term, it sequesters carbon of about 250 Mg C/ha. When a sole rubber plantation system is converted to rubber agroforestry system, the C sequestration potential of the plantation increases. In a study carried out in Indonesia by Palm et al., (1999), they found that carbon stocks increased from 46Mg C/ha to 89Mg C/ha when a 30 years rotational rubber were compared with permanent jungle-rubber. In a similar study in Indonesia, it was observed by that there was an increase in carbon stocks from 38Mg C/ha to 91Mg C/ha when a sole rubber system were converted to a rubber agroforestry system.

4. CONCLUSION

Rubber trees absorbed carbon dioxide in the atmosphere and stored it in its biomass, litter, and understory vegetation and in the soil. With this, rubber plantation can act as carbon sink. Rubber plantation have the potential to partially mitigate greenhouse gases by trapping carbon in the atmosphere. Carbon sequestration potential in rubber plantations is achieved by establishing rubber plantation on degraded forest or agricultural land. For example, the transformation of agrarian to grassland ecosystems is a common problem, and converting these land-use types into rubber plantations could sequester carbon and also generate income and foreign exchange from produce. However, converting mature forest to agrarian usually results to large carbon debts except when it is cultivated with rubber trees where the carbon payback time is around 40 years. Therefore, rubber plantations are established in low-carbon, severely degraded agricultural or forest land.

Ethical approval

Not applicable.

Informed consent

Not applicable.

Conflicts of interests

The authors declare that there are no conflicts of interests.

Funding

The study has not received any external funding.

Data and materials availability

All data associated with this study are present in the paper.

REFERENCES AND NOTES

- Blagodatsky S, Xu J, Cadisch G. Carbon balance of rubber (*Hevea brasiliensis*) plantations: a review of uncertainties at plot, landscape and production level. *Agric Ecosyst Environ* 2016; 221:8–19. doi: 10.1016/j.agee.2016.01.025
- Brahma B, Nath AJ, Sileshi GW, Das AK. Estimating biomass stocks and potential loss of biomass carbon through clear-felling of rubber plantations. *Biomass Bioenergy* 2018; 115:88–96. doi: 10.1016/j.biombioe.2018.04.019
- Cheng CM, Wang RS, Jiang JS. Variation of soil fertility and carbon sequestration by planting *Hevea brasiliensis* in Hainan Island, China. *J Environ Sci* 2007; 19(3):348–352. doi: 10.1016/S1001-0742(07)60057-6
- Choudhary BK, Majumdar K, Datta B. Carbon sequestration potential and edaphic properties along the plantation age of rubber in Tripura, Northeastern India. *Curr World Environ* 2016; 11:756–766. doi: 10.12944/CWE.11.3.10
- Day SK. A preliminary estimation of carbon stock sequestered through rubber (*Hevea brasiliensis*) plantation in north-eastern region of India. *Indian For* 2005; 131(11).
- FAO. Organic Agriculture, Environmental and Food Security. Environmental and Natural Resources Series, No 4, Rome, 2002.
- Fox J, Castella JC, Ziegler AD. Swidden, rubber and carbon: Can REDD+ work for people and the environment in Montane Mainland Southeast Asia? *Glob Environ Change* 2014; 29:318–326.
- Grieco E. Land use change and carbon stock dynamics in Sub-Saharan Africa Case study of Western Africa – Ghana. *Universit`a Degli Studi Della Tuscia, Italy*, 2011.
- Houghton RA, Van-der-Werf GR, DeFries RS, Hansen MC, House JI, Le-Quere C, Pongratz J, Ramankutty N. Chapter G2 Carbon emissions from land use and land cover change. *BG* 2012; 9:5125–5142. doi: 10.5194/bg-9-5125-2012
- IPCC. Intergovernmental Panel on Climate Change. IPCC guidelines for national greenhouse gas inventories. In: Eggleston H, Buendia L, Miwa K, Ngara T, Tanabe K (eds). IPCC-IGES, Japan, 2006.
- IRSG. Rubber Statistical Bulletin. Changi, Singapore, October–December: International Rubber Study Group (IRSG) 2019; 74 :4-6.
- Kiyono Y, Furuya N, Fujita N, Sato T, Matsumoto M, Bounthabandit S, Sanonty S. Can converting slash-and-burn agricultural fields into rubber tree (*Hevea brasiliensis*) plantations provide climate change mitigation? A case study in northern Laos. *FFPRI Bulletin* 2014; 13(3):79–88.
- Kongsager R, Napier J, Mertz O. The carbon sequestration potential of tree crop plantations. *Mitig Adapt Strateg Glob Change* 2013; 18:1197–1213. doi: 10.1007/s11027-012-9417-z
- Liguori G, Gugliuzza G, Inglese P. Evaluating carbon fluxes in orange orchards in relation to planting density. *J Agric Sci* 2009; 147(6):637–645. doi: 10.1017/S002185960900882X
- Liu C, Pang J, Jepsen M, Lü X, Tang J. Carbon stocks across a fifty-year chrono sequence of rubber plantations in tropical China. *Forests* 2017; 8(6):209. doi: 10.3390/f8060209
- Maggiotto SR, Oliveira DD, Marur CJ, Stivari SMS, Leclerc M, Wagner-Riddle C. Potential carbon sequestration in rubber tree plantations in the northwestern region of the Parana State, Brazil. *Acta Sci Agron* 2014; 36(2):239–245. doi: 10.4025/actasciagron.v36i2.17404
- Mesike CS, Sagay GA, Okoh RN. Optimum Replacement Period for Rubber Plantation in Nigeria. *Q J Int Agric* 2010; 49 (3):257–270.
- Nizami SM, Yiping Z, Liqing S, Zhao W, Zhang X. Managing Carbon Sinks in Rubber (*Hevea brasiliensis*) Plantation by Changing Rotation length in SW China. *PLoS One* 2014; 9(12): e11524. doi: 10.1371/journal.pone.0115234
- Palm CA, Woome PL, Alegre J, Arevalo L, Castilla C, Cordeiro DG, Feigl B, Hairiah K, Kotto-Same J, Mendes A, Moukam A, Murdiyars D, Njomgang R, Parton WJ, Ricse A, Rodrigues V, Sitompul SM, Van-Noordwijk M. Carbon Sequestration and Trace Gas Emissions in Slash-and-Burn and Alternative Land-Uses in the Humid Tropics. Nairobi, Kenya, ASB Climate Change Working Group Final Report, Phase II, ASB Coordination Office, ICRAF, Nairobi, Kenya, 1999.
- Petsri S, Chidthaisong A, Pumijumnong N, Wachrinrat C. Greenhouse gas emissions and carbon stock changes in rubber tree plantations in Thailand from 1990 to 2004. *J Clean Prod* 2013; 52:61–70. doi: 10.1016/j.jclepro.2013.02.003
- Song Q, Zhang Y. Biomass, carbon sequestration and its potential of rubber plantations in Xishuangbanna, Southwest China. *Chin J Ecol* 2010; 29(10):1887–1891.
- Soto-Pinto L, Anzueto M, Mendoza J, Ferrer GJ, De-Jon B. Carbon sequestration through agroforestry in indigenous

- communities of Chiapas, Mexico. *Agrofor Syst* 2010; 78:39–51. doi: 10.1007/s10457-009-9247-5
23. Sun Y, Ma Y, Cao K, Li H, Shen J, Liu W, Di L, Mei C. Temporal changes of ecosystem carbon stocks in rubber plantations in Xishuangbanna, Southwest China. *Pedosphere* 2017; 27(4):737–746. doi: 10.1016/S1002-0160(17)60327-8
24. Verchot L, Van-Noordwijk M, Kandji S, Tomich T, Ong C, Albrecht A, Mackensen J, Bantilan C, Anupama KV, Palm C. Climate change: linking adaptation and mitigation through agroforestry. *Mitig Adapt Strateg Glob Chang* 2007; 12:901–918.
25. Warren-Thomas EM, Edwards DP, Bebbler DP, Chhang P, Diment AN, Evans TD, Lambrick, FH, Maxwell JF, Nut M, O’Kelly HJ, Theilade I, Dolman PM. Protecting tropical forests from the rapid expansion of rubber using carbon payments. *Nat Commun* 2018; 9(1):911. doi: 10.1038/s41467-018-03287-9
26. Yang X, Blagodatsky S, Lippe M, Liu F, Hammond J, Xu J, Cadisch G. Land-use change impact on time-averaged carbon balances: Rubber expansion and reforestation in a biosphere reserve, South-West China. *For Ecol Manage* 2016; 372(3):149–163. doi: 10.1016/j.foreco.2016.04.009
27. Yang X, Blagodatsky S, Marohn C, Liu H, Golbon R, Xu J, Cadisch G. Climbing the mountain fast but smart: Modelling rubber tree growth and latex yield under climate change. *For Ecol Manage* 2019; 439:55–69.
28. Yang X, Blagodatsky S, Liu F, Beckschafer P, Xu J, Cadisch G. Rubber tree allometry, biomass partitioning and carbon stocks in mountainous landscapes of sub-tropical China. *For Ecol Manage* 2017; 404:84–99.
29. Ziegler AD, Fox JM, Xu J. The Rubber Juggernaut. *Sci* 2009; 324(5930):1024–1025. doi: 10.1126/science.1173833
30. Ziegler AD, Phelps J, Yuen JQ, Webb EL, Lawrence D, Fox JM, Bruun TB, Leisz SJ, Ryan CM, Dressler W, Mertz O, Pascual U, Padoch C, Koh LP. Carbon outcomes of major land-cover transitions in SE Asia: great uncertainties and gREDD plus policy implications. *Glob Change Biol* 2012; 18(10):3087–3099.